



# Effects of MHD on fast ion confinement in SPARC

RA Tinguely<sup>1</sup>, A Braun<sup>2</sup>, S Scott<sup>3</sup>, GJ Kramer<sup>4</sup>, M Podestà<sup>4</sup>

<sup>1</sup>*MIT PSFC*   <sup>2</sup>*PFURO*   <sup>3</sup>*CFS*   <sup>4</sup>*PPPL (INFUSE)*

This work was made possible by funding from the US Department of Energy for the Plasma and Fusion Undergraduate Research Opportunities (PFURO) program and the Innovation Network for Fusion Energy (INFUSE) initiative. This work is supported by the US DOE contract No. DE-AC02-09CH11466 and Commonwealth Fusion Systems.

# Outline

- Introducing SPARC
- Low frequency MHD, NTMs + TF ripple
- High frequency MHD, Alfven Eigenmodes
- Diagnostics, opportunities and challenges
- Summary and future work


# Introducing SPARC



# SPARC is a high-field, compact, DT tokamak under construction in Devens, MA; first plasma 2025

full-field DT H-mode	
$R_0$	1.85 m
$a$	0.57 m
$B_T$	12.2 T
$I_P$	8.7 MA
$q_{95}$	3.4
$\kappa_{95}$	1.75
$f_G$	0.37
$\beta_N$	1.0
$P_{ICRH}$	11.1 MW
$P_{fusion}$	140 MW
$Q$	11

[Creely 2020 JPP]




<https://www.psfc.mit.edu/sparc>

# SPARC will use only Ion Cyclotron Resonance Heating




- No Neutral Beam Injection
- $f = 120$  MHz at  $B_0 = 12$  T
  - Fundamental He3
  - Second harmonic T
- $P_{ICRH} \sim 25\text{-}30$  MW





[Lin 2020 JPP]

# SPARC's Primary Reference Discharge (PRD) has peak density $\sim 4 \times 10^{20} \text{ m}^{-3}$ and temperature $\sim 20 \text{ keV}$



[Creely 2020 JPP, Rodriguez Fernandez 2020 JPP]

# Fast ion (He3 minority) population is more peaked than alpha population in SPARC



[Scott 2020 JPP]

# SPARC and ITER have similar alpha parameters

- Alfven speed  $v_A = 9 \times 10^6 \text{ m/s}$
- Thermal D speed  $v_D = 1 \times 10^6 \text{ m/s}$
- Alpha speed  $v_\alpha = 13 \times 10^6 \text{ m/s}$

- Alpha Larmor radius  $\rho_\alpha \sim 2 \text{ cm}$
- Alpha banana orbit width  $\delta_\alpha < 12 \text{ cm}$

- Alpha beta  $\beta_\alpha \sim 0.6\%$
- Thermal beta  $\beta_{th} \sim 4\%$

- Unstable toroidal modes  $n = 13 - 20$
- High-B [Tolman 2019 NF]  $\rightarrow n = 30 - 40$

	ITER/SPARC	
	$Q = 3$	$Q = 9$
$\beta_\alpha / \beta$	2.46	1.23
$V_A$	0.94	0.97
$n_{\max}$	1.64	1.58



[Scott 2020 JPP]







# Low frequency MHD, NTMs + TF ripple

# Toroidal field ripple leads to alpha particle/ power loss ~ 500 kW



[Scott 2020 JPP]

# Neoclassical tearing modes $m/n = 3/2$ and $2/1$ modeled with SPIRAL




[Kramer 2013 PPCF, Braun 2022 In progress]

# Increase in alpha power loss is modest $\sim 2\% \rightarrow \sim 2.2\%$ , but total power loss remains low $< 700$ kW




Percentage of Total Alpha Power Lost by Ripple Case





[Braun 2022 In progress]

**First-wall heat flux from alphas increases by ~50%,  
~100 kW/m<sup>2</sup> → 150 kW/m<sup>2</sup> (flat wall assumption)**



[Braun 2022 In progress]

# Heat flux may be concentrated for a locked mode, but spread out with intrinsic rotation $<10$ kHz





- Upper-bound ripple
- Toroidally symmetric first wall

[Braun 2022 In progress]



# High frequency MHD, Alfven Eigenmodes

# $n = 10$ Alfvén continua and eigenmode structures calculated with NOVA (without rotation)





# $n = 15$ Alfvén continua and eigenmode structures calculated with NOVA (without rotation)




# NOVA-K calculates stability breakdown;


## He3 minority *stabilizes* AEs perhaps due to $\nu_{He3} \sim \nu_A$

Drive/Damping $\gamma/\omega_0$ (%)	n = 5 350 kHz	n = 10 482 kHz	n = 15 481 kHz	n = 20 370 kHz
Continuum	$\sim 0.0$	$\sim 0.0$	$\sim 0.0$	$\sim 0.0$
Radiative	-0.3	$\sim 0.0$	$\sim 0.0$	-2.6
Electron collisional	$\sim 0.0$	$\sim 0.0$	$\sim 0.0$	$\sim 0.0$
Electron Landau	-0.1	$\sim 0.0$	$\sim 0.0$	$\sim 0.0$
Ion Landau	$\sim 0.0$	$\sim 0.0$	$\sim 0.0$	-0.1
He3 minority	-0.1	-0.1	-0.1	-0.4
Alphas	$\sim 0.0$	+0.3	+0.3	+0.7
<i>Intrinsic damping</i>	-0.4	-0.1	-0.1	-2.8
<b>Total growth rate</b>	-0.5	+0.2	+0.1	-2.4

# ORBIT calculates wave-particle interactions over *all* of phase-space: $n = 10$ TAE + 3.5 MeV alphas



# ORBIT calculates wave-particle interactions over *all* of phase-space: $n = 10$ TAE + 2.4 MeV He3




# **Diagnostics, opportunities and challenges**



# At least one high-frequency Mirnov array is planned for AE measurements


- $n \leq 40$
- $m \leq 120$  (for  $q \leq 3$ )
- $f_{\text{TAE}} = \sim 500 \text{ kHz} + 40 \cdot 10 \text{ kHz} = 900 \text{ kHz}$
- $f_{\text{EAE}} = \sim 1 \text{ MHz} + 40 \cdot 10 \text{ kHz} = 1.4 \text{ MHz}$
- Digitization rate  $> 2 \text{ MS/s}$
- Other diagnostics:
  - Interferometry (density)
  - Soft X-Ray (temperature)



[R Sweeney]


# Neutron camera/spectrometer is planned for neutron/alpha birth profile measurements

- Neutron flux sufficiently high  
→ spectrometers along several lines-of-sight
  - Magnetic Proton Recoil
  - Compact CVD diamond
- High-energy neutron tail ( $<20$  MeV)
  - ICRH tail
  - Alpha knock-on neutrons




# Scoping underway for gamma ray diagnostics

- Look at *confined* alpha population
- Incorporate into neutron camera lines-of-sight?
- Reactions of interest:
  - No Beryllium (unlike JET and ITER)
  - Boronization:  
 $^{10}\text{B}(\alpha, \gamma)^{13}\text{C} \rightarrow \sim 3.1, 3.7, 3.85 \text{ MeV gammas}$
  - He3 minority heating:  
 $\alpha(^3\text{He}, \gamma)^7\text{Be} \rightarrow \sim 3\text{e}12 \text{ gammas/s in PRD}$
  - $\text{D}+\text{T} \rightarrow (1\text{e-}5) ^5\text{He}^* \rightarrow ^5\text{He} + \gamma (16.7 \text{ MeV})$



HPGe

LaBr<sub>3</sub>



[Tardocchi 2020 Varenna]



# Summary and future work



# Summary



- SPARC is under construction, with first plasma ~2025 and PRD ~2027
- TF ripple alone could cause alpha power loss  $<2.2\%$  ( $<620$  kW), and large  $3/2$  or  $2/1$  NTMs could increase this to  $<2.4\%$  ( $<680$  kW)
- $n = 10 - 15$  TAEs could be (marginally) destabilized by alphas, while He3 is interestingly stabilizing, near  $q = 1$  in the PRD
- Diagnostics for the alpha birth profile and high-frequency fluctuations are being designed

# Future work

- Evaluate other SPARC scenarios, e.g.  $Q \sim 1$  and 5
- Assess other low-frequency modes, e.g. sawteeth and ELMs
- Perform toroidal mode number scan and more advanced simulations
- Scope diagnostics for confined alpha particles
- Collaborations are very welcome!

# Bonus



# NOVA-K calculates stability breakdown;


## He3 minority *stabilizes* AEs perhaps due to $v_{He3} \sim v_A$

Drive/Damping $\gamma/\omega_0$ (%)	n = 5		n = 10		n = 15		n = 20	
	292 kHz	350 kHz	361 kHz	482 kHz	360 kHz	481 kHz	341 kHz	370 kHz
Continuum	−0.1	~0.0	~0.0	~0.0	~0.0	~0.0	−0.2	~0.0
Radiative	−2.7	−0.3	−1.4	~0.0	−1.0	~0.0	−6.4	−2.6
Electron collisional	~0.0	~0.0	~0.0	~0.0	~0.0	~0.0	~0.0	~0.0
Electron Landau	−0.1	−0.1	~0.0	~0.0	~0.0	~0.0	~0.0	~0.0
Ion Landau	~0.0	~0.0	−0.1	~0.0	−0.1	~0.0	~0.0	−0.1
He3 minority	−0.2	−0.1	−0.2	−0.1	−0.2	−0.1	−0.2	−0.4
Alphas	+0.1	~0.0	+0.6	+0.3	+0.8	+0.3	+0.2	+0.7
<i>Total w/o fast ions</i>	−2.9	−0.4	−1.6	−0.1	−1.1	−0.1	−6.6	−2.8
<b>Total w/ fast ions</b>	−3.0	−0.5	−1.1	+0.2	−0.5	+0.1	−6.6	−2.4

# ORBIT calculates wave-particle interactions ( $n = 10$ TAE + alphas) over *all* of phase-space

Trapped particles

Passing particles



wave  $\rightarrow$  particles

particles  $\rightarrow$  wave

# ORBIT calculates wave-particle interactions ( $n = 10$ TAE + He3) over *all* of phase-space

Trapped particles

Passing particles

